

Regulation of Gene Expression for L-Isoaspartyl O-Methyltransferases by Cis-Elements Associated with “Heat-Shock Polytene Chromosome Puffing Formation” in the Anhydrobiotic Midge

Deviatliarov R., Shagimardanova E., Kikawada T.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2016, Springer Science+Business Media New York. L-isoaspartyl O-methyltransferases (PIMTs) are known mostly because of their protein chain repair activity in plant seeds, but also in many other organisms, from bacteria to mammals, including human. The main function of PIMT is recognition and methylation of spontaneously isomerized aspartate residue (Asp) in the amino acid sequences of cellular proteins, and thus preventing their conformation shifts and subsequent inactivation or loss of particular functions. The correlation between PIMT activity and accumulation of isomerized aspartate residues is well studied and used as a marker of isoAsp presence, whereas regulation pathways of PIMT genes are not clear yet. In our study, we propose a possible model for multiple paralogous PIMT gene regulation in the genome of an anhydrobiotic insect *Polypedilum vanderplanki*. Analysis of promoter regions revealed that PIMT genes possess enriched motif nearby +1 site, associated with GO term heat-shock polytene chromosome puffing. Poly(ADP-ribose) polymerase (PARP) is involved in this process of chromatin loosening and subsequent transcription activation of related genes. This result implied that expression of PIMT genes in *P. vanderplanki* could be regulated by chromatin modification.

<http://dx.doi.org/10.1007/s12668-016-0312-z>

Keywords

PARP (poly(ADP-ribose) polymerase), PIMT (L-isoaspartyl O-methyltransferase), *Polypedilum vanderplanki*, Transcription regulation

References

- [1] Aswad, D., Paranandi, M., Schurter, B. (2000). Isoaspartate in peptides and proteins: formation, significance, and analysis. *Journal of Pharmaceutical and Biomedical Analysis*, 21, 1129–1136.
- [2] Clarke, S. (2003). Aging as war between chemical and biochemical processes: protein methylation and the recognition of age-damaged proteins for repair. *Ageing Research Reviews*, 2, 263–285.
- [3] Xu, Q., Belcastro, M., Villa, S., Dinkins, R., Clarke, S., Downie, A. (2004). A second protein L-isoaspartyl methyltransferase gene in *Arabidopsis* produces two transcripts whose products are sequestered in the nucleus. *Plant Physiology*, 136, 2652–2664.

- [4] Ogé, L., Bourdais, G., Bove, J., Collet, B., Godin, B., Granier, F., et al. (2008). Protein repair L-isoaspartyl methyltransferase 1 is involved in both seed longevity and germination vigor in *Arabidopsis*. *Plant Cell*, 20, 3022–3037.
- [5] Gusev, O., Suetsugu, Y., Cornette, R., Kawashima, T., Logacheva, M. D., Kondrashov, A. S., et al. (2014). Comparative genome sequencing reveals genomic signature of extreme desiccation tolerance in the anhydrobiotic midge. *Nature Communications*, 5, 4784. doi:10.1038/ncomms5784.
- [6] Nakahara, Y., Imanishi, S., Mitsumasu, K., Kanamori, Y., Iwata, K., Watanabe, M., et al. (2010). Cells from an anhydrobiotic chironomid survive almost complete desiccation. *Cryobiology*, 60, 138–146.
- [7] Kim, D., Pertea, G., Trapnell, C., Pimentel, H., Kelley, R., Salzberg, S. L. (2013). TopHat2: accurate alignment of transcriptomes in the presence of insertions, deletions and gene fusions. *Genome Biology*, 14, R36.
- [8] Watanabe, K., Imanishi, S., Akiduki, G., Cornette, R., Okuda, T. (2016). Air-dried cells from the anhydrobiotic insect, *Polypedilum vanderplanki*, can survive long term preservation at room temperature and retain proliferation potential after rehydration. *Cryobiology*, 73, 93–98.
- [9] Pirrotta, V. (2003). Transcription. Puffing with PARP. *Science*, 299, 528–529.
- [10] Tulin, A., & Spradling, A. (2003). Chromatin loosening by poly(ADP)-ribose polymerase (PARP) at *Drosophila* puff loci. *Science*, 299, 560–562.
- [11] Grube, K., & Bürkle, A. (1992). Poly(ADP-ribose) polymerase activity in mononuclear leukocytes of 13 mammalian species correlates with species-specific life span. *Proceedings of the National Academy of Sciences*, 89, 11759–11763.
- [12] Gusev, O., Nakahara, Y., Vanyagina, V., Malutina, L., Cornette, R., Sakashita, T., et al. (2010). Anhydrobiosis-associated nuclear DNA damage and repair in the sleeping chironomid: linkage with radioresistance. *PloS One*, 5, e14008.